



ELSEVIER

Cell biology of Ca^{2+} -triggered exocytosis

Zhiping P Pang¹ and Thomas C Südhof^{1,2}

Ca^{2+} triggers many forms of exocytosis in different types of eukaryotic cells, for example synaptic vesicle exocytosis in neurons, granule exocytosis in mast cells, and hormone exocytosis in endocrine cells. Work over the past two decades has shown that synaptotagmins function as the primary Ca^{2+} -sensors for most of these forms of exocytosis, and that synaptotagmins act via Ca^{2+} -dependent interactions with both the fusing phospholipid membranes and the membrane fusion machinery. However, some forms of Ca^{2+} -induced exocytosis may utilize other, as yet unidentified Ca^{2+} -sensors, for example, slow synaptic exocytosis mediating asynchronous neurotransmitter release. In the following overview, we will discuss the synaptotagmin-based mechanism of Ca^{2+} -triggered exocytosis in neurons and neuroendocrine cells, and its potential extension to other types of Ca^{2+} -stimulated exocytosis for which no synaptotagmin Ca^{2+} -sensor has been identified.

Addresses

¹ Department of Molecular and Cellular Physiology, Stanford University, 1050 Arastradero Rd., Palo Alto, CA 94304-5543, USA

² Howard Hughes Medical Institute, Stanford University, 1050 Arastradero Rd., Palo Alto, CA 94304-5543, USA

Corresponding authors: Pang, Zhiping P (zpang@stanford.edu) and Südhof, Thomas C (tcs1@stanford.edu)

Current Opinion in Cell Biology 2010, 22:496–505

This review comes from a themed issue on
Membranes and organelles
Edited by Suzanne Pfeffer and Peter Novick

Available online 3rd June 2010

0955-0674/\$ – see front matter

© 2010 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.ceb.2010.05.001

Introduction

Ca^{2+} -induced exocytosis initiates many forms of intercellular communication, as exemplified by synaptic transmission, which begins with Ca^{2+} -triggered synaptic vesicle exocytosis that mediates neurotransmitter release (Figure 1) [1]. Similarly, neuroendocrine cells secrete hormones by Ca^{2+} -induced exocytosis [2], mast cells release their granule contents upon stimulation by Ca^{2+} -controlled exocytosis [3], and even in T-lymphocytes, Ca^{2+} -triggered exocytosis is functionally essential [4]. The question of how Ca^{2+} triggers exocytosis was first raised by Bernhard Katz's seminal discovery that Ca^{2+} induces synaptic vesicle exocytosis, and thereby initiates synaptic transmission [5]. However, not much progress

was made in this question until the discovery of synaptotagmin-1 (Syt1) as a candidate Ca^{2+} -sensor for synaptic exocytosis [6]. Work from many laboratories has provided overwhelming evidence that Syt1 and its homologs function as the primary Ca^{2+} -sensors in most forms of exocytosis, and has elucidated the principal mechanism by which synaptotagmin operates [1]. However, as described below, this work has also raised important new questions about the role of Ca^{2+} in regulating membrane traffic. The present review focuses on the cell biology of Ca^{2+} -triggered exocytosis in neurons and endocrine cells, and tries to relate the emerging synaptotagmin Ca^{2+} -sensor paradigm to these new unanswered questions.


Synaptic exocytosis

In presynaptic nerve terminals, neurotransmitters are packaged into small synaptic vesicles, and released by Ca^{2+} -triggered exocytosis of synaptic vesicles at the presynaptic active zone (Figure 1). Three different modes of neurotransmitter release exist (Figure 2):

1. Evoked synchronous release initiates within a millisecond after an action potential induces Ca^{2+} -influx into a presynaptic terminal [7,8]. Fast synchronous release measured as the postsynaptic response can be fitted by a double exponential function, and thus can be arbitrarily subdivided into a fast and a slow phase [9].
2. Evoked asynchronous release sets in with a delay after an action potential, and is normally negligible [10], either because it is outcompeted by the synchronous release mechanism [11], or because the synchronous release mechanism (i.e. synaptotagmin and complexin, see below) suppresses asynchronous release [12]. However, asynchronous release becomes a dominant form of release in some synapses during high-frequency trains of action potentials, particularly in inhibitory synapses [13^{••},14^{••},15].
3. Spontaneous 'mini' release, finally, represents the exocytosis of single vesicles [16] that is independent of action potentials, but nevertheless largely Ca^{2+} -dependent [17^{••}]. Spontaneous release is induced by resting Ca^{2+} -concentrations, or may be stimulated by stochastic Ca^{2+} -channel openings and/or Ca^{2+} -sparks via Ca^{2+} -influx from internal Ca^{2+} -stores [18^{••}].

All three forms of synaptic exocytosis probably perform important physiological functions. Evoked synchronous release represents the primary mode of intercellular communication between neurons, rendering its importance obvious. Bursts of activity are often seen in neurons, suggesting that these trigger asynchronous release that

Figure 1




Synaptic and endocrine Ca²⁺-triggered exocytosis. At a synapse (**left**), neurotransmitters are packaged into small synaptic vesicles, which are docked at the active zone adjacent to voltage-dependent Ca²⁺-channels. A presynaptic action potential (*insert*) gates Ca²⁺-influx into the terminal, thereby triggering vesicle exocytosis. The released transmitters produce a postsynaptic current (*insert*) that can be recorded by whole-cell patch clamping. In endocrine cells (**right**), hormones are packaged into LDCVs, which are generally not docked. Upon sustained increases in cytosolic Ca²⁺, as obtained during stimulation or Ca²⁺-uncaging (*insert*), exocytosis is triggered with a significantly slower time course than at a synapse, as measured by amperometry or capacitance (C_m; *insert*). Note that Ca²⁺-channel and release sites are not tightly coupled in endocrine cells. ER, endoplasmic reticulum; M, mitochondrion. Traces are shown purely for demonstration purposes.

may shape neural function at the circuit level [19^{*}]. Spontaneous release may improve the signal to noise ratio at the systems level [20], and/or to contribute to the development and maintenance synapses [21].

In central synapses, synaptic vesicles exist in functionally different pools [22], although this may not be true for neuromuscular synapses [23]. All types of synaptic exocytosis probably utilize the same release-ready pool of vesicles (the so-called readily releasable pool [RRP]),

which accounts only for a small percentage of the total vesicle pool [22]. An operational definition of the RRP is the amount of release triggered by application of hypertonic sucrose, which induces Ca²⁺-independent exocytosis of all synaptic vesicles in the RRP [24,25]. The morphological correlate of the RRP is unclear; RRP vesicles are distinct from the synaptic vesicles that are morphologically docked at the active zone because the number of docked vesicles does not correlate with the RRP size [26], and a point mutation that activates the

Figure 2



Synaptic vesicle exocytosis detected by whole-cell patch clamp recordings. Images depict representative traces of postsynaptic currents illustrating the three different forms of synaptic exocytosis: evoked synchronous release from wild-type synapses (**a**), evoked asynchronous release from Syt1-deficient synapses (**b**) and spontaneous 'mini' release (**c**). Note that asynchronous release also can be recorded in some wild-type neurons upon high-frequency stimulation.

SNARE-protein syntaxin-1B simultaneously increases the number of docked vesicles but decreases the RRP size [27**].

Endocrine exocytosis

Hormonal exocytosis of endocrine cells operates on large dense-core vesicles (LDCVs) that are probably similar to neuropeptide LDCVs in neurons. LDCV exocytosis has been studied mostly in adrenal chromaffin cells and in pancreatic β -cells, where Ca^{2+} -triggered exocytosis operates in three phases, referred to as the fast, slow, and sustained phase [28,29]. The three phases of LDCV exocytosis have been attributed to different vesicle pools, but it is uncertain whether these pools represent physically distinct types of vesicles, or simply different functional states. In addition, spontaneous exocytosis of LDCVs has also been described in chromaffin cells, and also appears to be regulated by Ca^{2+} [30].

Even the fast phase of LDCV exocytosis is much slower than both synchronous and asynchronous synaptic exocytosis (Figure 1). Nevertheless, LDCV and synaptic exocytosis are very similar, as they appear to use the same Ca^{2+} -triggering mechanisms (see discussion below), and differ primarily in how synaptic vesicles and LDCVs are docked and prepared for fusion (i.e. primed).

Synaptotagmins as Ca^{2+} -sensors for exocytosis

Synaptotagmins are synaptic and secretory vesicle proteins (although some isoforms may be on the plasma membrane) that contain a single N-terminal transmembrane region, and two C-terminal Ca^{2+} -binding C2-domains [6]. 16 mammalian synaptotagmin isoforms were identified, 8 of which bind Ca^{2+} with distinct apparent Ca^{2+} -affinities (Syt1-3, Syt5-7, Syt9, and Syt10 [31–33], Table 1). Synaptotagmins are highly conserved evolutionarily; even all invertebrates express multiple isoforms. However, synaptotagmins are absent from plants and unicellular eukaryotes, suggesting that they emerged coincidentally with animals during evolution.

Synaptic and endocrine exocytosis are mediated by the same fusion machinery composed of SNARE-proteins and SM-proteins as other membrane fusion reactions—in fact, this fusion machinery was discovered at the synapse [34]. SNARE-proteins catalyze fusion by forming a complex that bridges the two fusing membranes, forcing these membranes together, whereas SM-proteins promote fusion by an unknown but essential mechanism. Ca^{2+} -binding to synaptotagmin triggers exocytosis by operating on this fusion machinery with the help of an ancillary protein called complexin. Thus, only six proteins – three SNARE-proteins, one SM-protein (Munc18-1), one synaptotagmin, and complexin – form the core of the Ca^{2+} -triggered exocytosis machinery (Figure 3), constituting a molecular clockwork that

exhibits an amazing simplicity that we refer to as the synaptotagmin paradigm [34].

Both C2-domains of synaptotagmin bind Ca^{2+} ; C2-domains were first shown to represent Ca^{2+} -binding domains in Syt1 [35]. C2-domains are janus-faced domains with 2-3 Ca^{2+} -binding sites on top, and a Ca^{2+} -independent surface on the bottom. Ca^{2+} -binding induces simultaneous synaptotagmin-binding to both the fusing phospholipid membranes, and the assembling SNARE complex (Figure 3). Complexin activates SNARE-complexes before synaptotagmin action, and clamps fusion by preventing complete SNARE-complex assembly until Ca^{2+} binds to synaptotagmin [36**,37**]. Complexin performs these actions by binding to SNARE complexes, and synaptotagmin dislodges the complexin clamp (Figure 3 step 5, [38]). Blocking Ca^{2+} -binding to the C2B-domain blocks synchronous exocytosis [39], whereas blocking Ca^{2+} -binding to the C2A-domain decreases exocytosis ~40%, and additionally decreases the apparent Ca^{2+} -cooperativity of exocytosis ~40% [40*]. Thus, the two C2-domains of synaptotagmins thus are not equivalent, but they cooperate with each other, with the C2B-domain playing the leading part. Nevertheless, mutations in the C2A-domain alter the overall apparent Ca^{2+} -affinity of synaptotagmin and change the Ca^{2+} -affinity of synaptic exocytosis correspondingly, an observation that provided the formal proof for the Ca^{2+} -sensor of synaptotagmin in exocytosis [41,42]. Biochemically, Ca^{2+} -binding to the C2B-domain is essential both for effective Ca^{2+} -dependent phospholipid binding of synaptotagmin and for displacing the complexin clamp from SNARE complexes. Note that in the latter process, complexin may still remain associated with the SNARE complex, since it probably interacts with the complex via multiple mechanisms.

Three synaptotagmins (Syt1, Syt2, and Syt9) function as Ca^{2+} -sensors for synaptic exocytosis, but exhibit distinct expression patterns and properties [43*]. Syt2 triggers release much faster than Syt1 and operates in synapses relying on fast signaling, such as the calyx of Held synapse or the neuromuscular junction [8–10,44**]. Syt9 triggers release with a significantly slower time course than either Syt1 or Syt2, and is primarily expressed in the reward pathway [43*].

In endocrine LDCV exocytosis, the major Ca^{2+} -sensors are Syt1 and Syt7, another Ca^{2+} -binding synaptotagmin [45–47,48*,49*,50*]. Single deletions of Syt1 or Syt7 in chromaffin cells impair preferentially the fast or slow phase of LDCV exocytosis, respectively, suggesting that the two synaptotagmins mediate different phases, whereas double deletions of Syt1 and Syt7 block both phases [48*]. Point mutations in Syt1 that change its Ca^{2+} -affinity change the apparent Ca^{2+} -affinity of exocytosis in chromaffin cells correspondingly, similar to the synapse [46]. Whereas at least Syt2 can substitute for Syt1

Table 1

Properties of synaptotagmins.

Protein	Class	Localization	Ca ²⁺ -binding	Special properties	Function	References
Syt 1	A	Synaptic vesicles LDCVs	Yes	N-terminal N-glycosylation Phospholipids and SNARE binding	Ca ²⁺ -sensor for fast exocytosis	[6,10,12,31–33,39, 40*,41,42,43*,74]
Syt2	A	Synaptic vesicles LDCVs?	Yes	N-terminal N-glycosylation Phospholipid and SNARE binding	Ca ²⁺ -sensor for fast exocytosis	[8,9,31–33,43*, 44**,54*,74]
Syt3	C	Plasma membrane	Yes	Disulfide bonds at N-terminus Phospholipid binding	Ca ²⁺ -sensor for exocytosis?	[31–33,74]
Syt4	D	Postsynaptic?	No*	Aspartate to Serine substitution in C2A domain loops	Unknown [†] ; not essential for survival	[73,74]
Syt5	C	Plasma membrane	Yes	Disulfide bonds at N-terminus Phospholipid bindings	Ca ²⁺ -sensor for exocytosis?	[31–33,74]
Syt6	C	Plasma membrane	Yes	Disulfide bonds at N-terminus Phospholipid binding	Ca ²⁺ -sensor for exocytosis?	[31,74]
Syt7	B	LDCVs Synapses?	Yes	Multiple splicing isoforms Phospholipid and SNARE binding	Ca ²⁺ -sensor for LDCV exocytosis	[47,48*,49*,50*,74]
Syt8	F	Primarily GIT tract	No	Only Syt isoform that is expressed at highest levels outside of brain	Unknown	[31,74]
Syt9	A	Synaptic vesicles	Yes	Phospholipid and SNARE binding	Ca ²⁺ -sensor for fast exocytosis	[43*,74]
Syt10	C	Plasma membrane	Yes	Disulfide bonds at N-terminus Phospholipid binding	Ca ²⁺ -sensor for exocytosis?	[31,74]
Syt11	D	Unknown	No	Similar to Syt4, but more abundant	Unknown	[74]
Syt12	F	Synaptic vesicles	No	Phosphorylated by PKA	May regulate miniature release	[31,74,75]
Syt13	F	Unknown	No		Unknown	[31,74]
Syt14	E	Unknown	No	Syt14-16 form a group of related and evolutionarily conserved isoforms	Unknown	[76]
Syt15	E	Unknown	No	Syt14-16 form a group of related and evolutionarily conserved isoforms	Unknown	[76]
Syt16	E	Unknown	No	Syt14-16 form a group of related and evolutionarily conserved isoforms	Unknown	[76]

Synaptotagmins are defined by the presence of an N-terminal transmembrane region, and two C-terminal C2-domains. Each synaptotagmin protein is encoded by a different gene; some are differentially spliced (e.g. Syt7). Classes of synaptotagmins are proposed defined by sequence homologies, conservation, and functions. A further isoform, 'synaptotagmin-17', is not listed because it lacks a transmembrane region, and thus is not a bona fide synaptotagmin.

* Syt4 binds to Ca²⁺ in *Drosophila* and has been implicated in postsynaptic exocytosis.

in chromaffin exocytosis [51], Syt7 cannot substitute for Syt1 in synaptic exocytosis [43*], and Syt7 deletions have no significant effect on synaptic exocytosis [52]. The reason for this selectivity remains unclear, as Syt7 is very similar to Syt1, and only differs from Syt1 in lacking an N-terminal intravesicular N-glycosylation site and in exhibiting a higher apparent Ca²⁺-affinity [32]. In chromaffin cells, Syt1 may also be involved in docking of LDCVs since LDCV docking was impaired in cells from Syt1 KO mice [53*]. However, the mechanism of this change remains unclear because synapses lacking Syt1 exhibit no major docking phenotype, and a similar docking phenotype was observed in chromaffin cells but not synapses in other mutant mice [27**].


Synaptotagmins have also been implicated in non-neuronal and non-endocrine forms of Ca²⁺-induced exocytosis. In mast cells, Syt2 functions as the major Ca²⁺-sensor for exocytosis [54*]. In fibroblasts, Syt7

may mediate Ca²⁺-triggered lysosome exocytosis [55], although this has been disputed [56]. It is notable that no definitive functions were described for 4 of the 8 Ca²⁺-binding synaptotagmin isoforms (i.e. Syt3, Syt5, Syt6, and Syt10). These synaptotagmins form a highly homologous subgroup that exhibit Ca²⁺-dependent phospholipid-binding and SNARE-binding similar to Syt1, Syt2, Syt7, and Syt9, and probably also function in exocytosis.

Dual-Ca²⁺-sensor model for synaptic exocytosis

The most precise definition of synaptic transmission was achieved at the giant calyx of Held synapse in the brainstem that allows simultaneous patching of presynaptic and postsynaptic cells [57,58]. Measurements in the calyx synapse provided estimates of the Ca²⁺-affinity (10–100 μM) and Ca²⁺-cooperativity (~5 Ca²⁺-ions) of neurotransmitter release [44**,57,58]. As shown in mutant mice, this release is mediated by Syt2 as Ca²⁺-sensor [9,44**].

Figure 3



Model of the molecular steps mediated synaptic vesicle exocytosis (modified from [38]). Synaptic vesicles are docked at the active zone of a presynaptic terminal with unassembled SNARE complexes (top) and are then primed for release by partial SNARE-complex assembly that is catalyzed by Munc18, Munc13, and RIM (step 1). At least in inhibitory synapses, this priming process might be further modulated by ELKS2. The primed vesicles form the substrate for two main pathways of Ca^{2+} -triggered neurotransmitter release: asynchronous release (steps 2 and 3), in which full assembly of SNARE complexes leads to fusion-pore opening followed by complete fusion (step 3); and synchronous release (steps 4, 5, and 6), in which 'superpriming' by binding of complexins to assembled SNARE complexes (step 4) activates and freezes SNARE complexes in a metastable state (referred to as priming stage II). This stage is then substrate for fast Ca^{2+} -triggering of release when Ca^{2+} -binding to Syt1 induces its binding to phospholipids and to SNARE complexes, with the latter reaction displacing complexin and resulting in fusion-pore opening (step 5) and full fusion (step 6). Both the synchronous and the asynchronous release pathway can mediate spontaneous 'mini' release, depending on the local Ca^{2+} -microdomain. Synaptotagmin and complexin clamp (**block, in red**) the unidentified slow Ca^{2+} -sensor that mediates the asynchronous release; this clamping is relieved when Ca^{2+} binds to Syt1, allowing competition between Syt1 and the asynchronous Ca^{2+} -sensor during high-frequency stimulation [12,44**].

Calyx synapses normally exhibit little asynchronous release, and produce almost only synchronous release even at stimulation frequencies of 100 Hz, making it impossible to analyze asynchronous release biophysically. In Syt2-deficient synapses, however, asynchronous release could be analyzed in isolation, uncontaminated by a more dominant synchronous release component [44**]. Such analyses uncovered an asynchronous component of exocytosis in calyx synapses that displayed an apparent Ca^{2+} -cooperativity of exocytosis of only ~ 2 Ca^{2+} -ions, whereas synchronous release operated with an apparent Ca^{2+} -cooperativity of ~ 5 Ca^{2+} -ions, although both exhibited similar Ca^{2+} -affinities (~ 40 μM). Because of the different Ca^{2+} -cooperativities but similar Ca^{2+} -affinities of synchronous and asynchronous exocytosis,

low Ca^{2+} -concentrations preferentially but incompletely activate asynchronous release in wild-type synapses, whereas higher Ca^{2+} -concentrations, such as those achieved following an action potential, preferentially and completely activate synchronous release [44**].

Based on the biophysical definition of asynchronous release, a dual- Ca^{2+} -sensor model was proposed that at present provides the most precise description of evoked synaptic vesicle exocytosis for synapses [44**]. The model assumes that the synchronous Ca^{2+} -sensor synaptotagmin competes with an unknown asynchronous Ca^{2+} -sensor, with the asynchronous Ca^{2+} -sensor binding Ca^{2+} more slowly but at lower concentrations than the synchronous Ca^{2+} -sensor. It should be noted that although the

dual-Ca²⁺-sensor model is the best available, it does not take into account the intrinsic heterogeneity of synapse vesicles. Vesicles probably differ in their proximity to Ca²⁺-channels, which is a major determinant for the probability and speed of exocytosis [59]. Moreover, the number of SNARE complexes on a primed vesicle probably contributes to the Ca²⁺-sensitivity of this vesicle, but differs between vesicles [27**]. Finally, synaptic vesicles exhibit size heterogeneity [60], resulting in variations in the postsynaptic signal. In addition, the dual-Ca²⁺-sensor model also does not include the possibility that at least at some synapses, synaptotagmin may inhibit the asynchronous Ca²⁺-sensor [12,17**]. Thus, despite the fact that the dual-Ca²⁺-sensor model is currently the best available, it is far from perfect.

Synaptotagmin as a Ca²⁺-sensor for spontaneous release

At a synapse, lowering the extracellular Ca²⁺-concentration partially blocks spontaneous mini release; incubating synapses with membrane-permeable Ca²⁺-buffers, however, or infusing Ca²⁺-buffers into the calyx presynaptic terminal, blocks almost all spontaneous release [9,17**]. These results suggested that the majority of spontaneous release is Ca²⁺-dependent, but raised the question what Ca²⁺-sensor mediates this effect. Interestingly, knockin mutations in Syt1 that change its apparent Ca²⁺-affinity caused corresponding effects on the frequency of spontaneous release. Specifically, when the apparent Ca²⁺-affinity of Syt1 is decreased ~2-fold by the R233Q mutation, the frequency of spontaneous release is decreased ~2-fold, whereas an increase in Ca²⁺-dependent SNARE-complex binding by the D232N mutation of Syt1 caused a correspondingly large increase in spontaneous release frequency [17**]. These effects suggested that most spontaneous release is induced by Ca²⁺-binding to Syt1, with the Ca²⁺ derived from resting Ca²⁺-levels, Ca²⁺-influx via stochastically opening Ca²⁺-channels, or Ca²⁺-sparks. However, a recent study showed that deletion of another Ca²⁺-binding protein, Doc2, causes a partial decrease in mini frequency, indicating that this protein may contribute to the Ca²⁺-regulation of spontaneous release [61*].

Strikingly, although point mutations in Syt1 modulate spontaneous release, deletion of Syt1 or Syt2 increases spontaneous release dramatically, despite blocking evoked synchronous release [9,12,44**]. The 'new' spontaneous release in Syt1-deficient synapses is still Ca²⁺-dependent, but activated at lower extracellular Ca²⁺-concentrations with a lower apparent Ca²⁺-cooperativity than wild-type spontaneous release [17**]. Thus, the increased spontaneous release in Syt1 KO synapses (or, for that matter, also in Syt2 KO synapses) exhibits the properties of asynchronous release as determined in the calyx synapse [44**]. These results indicate Syt1 and Syt2 might generally inhibit asynchronous release.

An alternative explanation is that spontaneous release represents a completely separate cell-biological pathway, as suggested by reports that evoked and spontaneous synaptic release are using different synaptic vesicle pool [62,63**]. If so, the Syt1 and Syt2 deletions may not actually disinhibit the asynchronous Ca²⁺-sensor, but instead activate this separate pathway.


Other potential Ca²⁺-sensors for exocytosis

Which Ca²⁺-sensor mediates asynchronous release and other Ca²⁺-dependent types of exocytosis in which Syt1, Syt2, Syt7, and Syt9 do not act as Ca²⁺-sensors? Naturally, prime candidates are the other four Ca²⁺-binding synaptotagmins that have no known function (Syt3, Syt5, Syt6, and Syt10). However, at least for asynchronous release, this candidacy is doubtful since these synaptotagmins bind Ca²⁺ via a mechanism akin to that of Syt1, with a likely Ca²⁺-binding stoichiometry of ~5, whereas asynchronous release exhibits a much lower apparent Ca²⁺-stoichiometry.

Two major classes of cytosolic Ca²⁺-binding proteins are known, EF-hand proteins and C2-domain containing proteins. Calmodulin, the most important EF-hand Ca²⁺-binding protein, appears to enhance neurotransmitter release in both excitatory and inhibitory synapses without directly participating in asynchronous exocytosis [64]. Numerous other EF-hand Ca²⁺-binding proteins are also expressed in brain, but most function as Ca²⁺-buffers or as Ca²⁺-regulated enzymes. Many C2-domain proteins are expressed in brain, most of which contain a single C2-domain and are involved in signal transduction. A smaller subset of C2-domain proteins contains multiple C2-domains, such as synaptotagmins (Figure 4). Although the function of most of these multiple C2-domain proteins is unknown, the synaptotagmin paradigm suggests that at least some of them are involved in membrane traffic, rendering them candidates for exocytotic Ca²⁺-sensors.

1. *Ferlins* contain at least six C2-domains and a C-terminal transmembrane region; the two C-terminal C2-domains include canonical Ca²⁺-binding sequences. Ferlins are required for sperm exocytosis in *C. elegans* [65]. Of the 5 mammalian ferlins, dysferlin is involved in Ca²⁺-dependent exocytosis of repair vesicles in muscle [66], and otoferlin is required for Ca²⁺-dependent exocytosis in hair cells [67], although the mechanisms involved have not been defined.
2. *MCTPs* (multiple C2-domain transmembrane proteins) are evolutionarily conserved proteins with 3 C2-domains containing canonical Ca²⁺-binding sequences, and a C-terminal transmembrane region. The functions of MCTPs are unknown [68].
3. *E-Syts* (extended synaptotagmins) are also evolutionarily conserved transmembrane proteins. E-Syts contain an N-terminal transmembrane region like

Figure 4



Structures of multiple C₂-domain proteins. **(a)** Schematic diagram of the top Ca^{2+} -binding loops of the Syt1 C₂A-domain (modified from [41]). The C₂A-domain has three top loops, of which loops 1 and 3 form three Ca^{2+} -binding sites. Five aspartate residues, one serine residue and two backbone carbonyl groups coordinate the three bound Ca^{2+} ions. Residues are shown in single-letter amino acid code and are identified by residue numbers corresponding to mouse Syt1. **(b)** Domain structures of proteins with multiple C₂-domains that contain or lack a transmembrane domain. The mammalian genome encodes four classes of multiple C₂-domain proteins containing transmembrane regions: synaptotagmins (Syt1-16; note that 'Syt17' is not listed because it does not contain a transmembrane region, but is membrane-anchored via palmitoylated cysteine residues), extended synaptotagmins (E-Syt1-3), multiple C₂-domain and transmembrane proteins (MCTP1-2) and ferlins (including otoferlin, dysferlin, myoferlin and 2 other uncharacterized ferlin-like proteins). Examples of soluble C₂-domain containing proteins are also included. C₂-domains with canonical Ca^{2+} -binding consensus sequences are labeled as Ca^{2+} -binding C₂-domains, although Ca^{2+} -binding has not yet been tested to all of these domains. C₂-domains were assigned based on the conserved domain database of the NCBI; some proteins, especially ferlins, may have additional unpredicted C₂-domains that do not precisely fit the consensus sequence, as well as alternative transcripts with fewer C₂-domains.

synaptotagmins, followed by a single SMP-domain and either five C₂-domains (E-Syt1) or three C₂-domains (E-Syt2 and E-Syt3). Only the N-terminal C₂-domain contains the requisite Ca^{2+} -binding sequence, and exhibits Ca^{2+} -dependent phospholipid binding [69]. No functional data on E-Syts are available.

4. *Synaptotagmin-like proteins (SLPs)* represent a large and heterogeneous class of proteins with two C-terminal C₂-domains but without transmembrane regions. An evolutionarily conserved subset of SLPs contains N-terminal Rab-binding sequences and zinc-finger domains (including rabphilin); another subset of SLPs that is not evolutionarily conserved lacks the N-terminal Zinc-finger domain (Doc2s, SLP2c, SLP3b). In vertebrate synapses, rabphilin is important for re-priming of synaptic vesicles [70]. Doc2 proteins appear to bind SNARE proteins and phospholipids tighter than Syt1, and deletion of Doc2 α and Doc2 β partly reduces spontaneous release [61]. In chromaffin cells, Doc2 β plays a role in Ca^{2+} -dependent priming and exocytosis [71], but the overall function of Doc2 proteins and other SLPs remains unclear.
5. *Copines* are soluble double C₂-domain proteins of unknown function that bind to phospholipids [72]. Copines are evolutionarily conserved, and contain two N-terminal C₂-domains with canonical Ca^{2+} -binding sequences and a unique C-terminal domain.

Conclusions

A universal mechanism by which Ca^{2+} -binding to synaptotagmins triggers exocytosis has emerged over the past decade. This mechanism mediates most Ca^{2+} -triggered exocytosis using a pas-de-deux of synaptotagmins and complexin acting on SNARE complexes and phospholipid membranes. However, new intriguing questions have emerged. Are forms of exocytosis for which no synaptotagmin Ca^{2+} -sensor has been identified, such as asynchronous release, mediated by an atypical synaptotagmin, or by novel Ca^{2+} -sensor, for example one of the other multiple C₂-domain proteins (Figure 4)? How do the Ca^{2+} -sensors for synchronous and asynchronous release intersect in synaptic exocytosis—do they compete, or do synaptotagmins and complexin inhibit the alternative pathway? What is the role of ferlins in exocytosis in Ca^{2+} -triggered exocytosis? Answering these questions will significantly advance the field beyond the synaptotagmin paradigm.

Note Added in Proof

After this review was accepted for publication, three more papers described single molecule studies demonstrating how synaptotagmin functions as a Ca^{2+} -sensor for Ca^{2+} -dependent vesicle fusion in vitro [77–79].

Acknowledgements

We wish to thank Drs. Weiping Han and Jianyuan Sun for valuable discussions. Z.P.P. was supported by NARSAD Young Investigator Award and NIH/NINDS Epilepsy Training Grant 5T32NS007280.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Südhof TC: **The synaptic vesicle cycle.** *Annu Rev Neurosci* 2004, **27**:509-547.
 2. Bean AJ, Zhang X, Hokfelt T: **Peptide secretion: what do we know?** *FASEB J* 1994, **8**:630-638.
 3. Lindau M, Gomperts BD: **Techniques and concepts in exocytosis: focus on mast cells.** *Biochim Biophys Acta* 1991, **1071**:429-471.
 4. Randriamampita C, Trautmann A: **Ca²⁺ signals and T lymphocytes; "New mechanisms and functions in Ca²⁺ signalling".** *Biol Cell* 2004, **96**:69-78.
 5. Katz B, Miledi R: **Ionic requirements of synaptic transmitter release.** *Nature* 1967, **215**:651.
 6. Perin MS, Fried VA, Mignery GA, Jahn R, Südhof TC: **Phospholipid binding by a synaptic vesicle protein homologous to the regulatory region of protein kinase C.** *Nature* 1990, **345**:260-263.
 7. Borst JG, Helmchen F, Sakmann B: **Pre- and postsynaptic whole-cell recordings in the medial nucleus of the trapezoid body of the rat.** *J Physiol* 1995, **489**:825-840.
 8. Pang ZP, Melicoff E, Padgett D, Liu Y, Teich AF, Dickey BF, Lin W, Adachi R, Südhof TC: **Synaptotagmin-2 is essential for survival and contributes to Ca²⁺ triggering of neurotransmitter release in central and neuromuscular synapses.** *J Neurosci* 2006, **26**:13493-13504.
 9. Pang ZP, Sun J, Rizo J, Maximov A, Südhof TC: **Genetic analysis of synaptotagmin 2 in spontaneous and Ca²⁺-triggered neurotransmitter release.** *EMBO J* 2006, **25**:2039-2050.
 10. Geppert M, Goda Y, Hammer RE, Li C, Rosahl TW, Stevens CF, Südhof TC: **Synaptotagmin I: a major Ca²⁺ sensor for transmitter release at a central synapse.** *Cell* 1994, **79**:717-727.
 11. Otsu Y, Shahrezaei V, Li B, Raymond LA, Delaney KR, Murphy TH: **Competition between phasic and asynchronous release for recovered synaptic vesicles at developing hippocampal autaptic synapses.** *J Neurosci* 2004, **24**:420-433.
 12. Maximov A, Südhof TC: **Autonomous function of synaptotagmin 1 in triggering synchronous release independent of asynchronous release.** *Neuron* 2005, **48**:547-554.
 13. Best AR, Regehr WG: **Inhibitory regulation of electrically coupled neurons in the inferior olive is mediated by asynchronous release of GABA.** *Neuron* 2009, **62**:555-565.
This paper shows that the GABAergic input into inferior olive neurons from deep cerebellar nuclei was almost exclusively asynchronous, with little conventional synchronous release. Stimulation of above 10 Hz evoked asynchronous steady-state inhibitory current that depends on residual calcium signaling.
 14. Daw MI, Tricoire L, Erdelyi F, Szabo G, McBain CJ: **Asynchronous transmitter release from cholecystokinin-containing inhibitory interneurons is widespread and target-cell independent.** *J Neurosci* 2009, **29**:11112-11122.
This study examined asynchronous release from cholecystokinin-containing hippocampal interneurons during trains of action potentials. The authors also studied the Ca²⁺-sensitivity of synchronous and asynchronous release from this synapse, and suggested that these two types of vesicle exocytosis are probably operated by different Ca²⁺-sensors.
 15. Hefft S, Jonas P: **Asynchronous GABA release generates long-lasting inhibition at a hippocampal interneuron-principal neuron synapse.** *Nat Neurosci* 2005, **8**:1319-1328.
 16. Katz B, Miledi R: **Tetrodotoxin and neuromuscular transmission.** *Proc R Soc Lond B Biol Sci* 1967, **167**:8-22.
 17. Xu J, Pang ZP, Shin OH, Südhof TC: **Synaptotagmin-1 functions as a Ca²⁺ sensor for spontaneous release.** *Nat Neurosci* 2009, **12**:759-766.
 18. Llano I, González J, Caputo C, Lai FA, Blayney LM, Tan YP, Marty A: **Presynaptic calcium stores underlie large-amplitude miniature IPSCs and spontaneous calcium transients.** *Nat Neurosci* 2009, **3**:1256-1265.
This paper reveals that intracellular presynaptic Ca²⁺-stores can drive spontaneous neurotransmitter release by generating spontaneous Ca²⁺-transients.
 19. Harvey CD, Collman F, Dombeck DA, Tank DW: **Intracellular dynamics of hippocampal place cells during virtual navigation.** *Nature* 2009, **461**:941-946.
This study uses an interesting experimental paradigm in mice, namely virtual navigation, in conjunction with electrophysiology to describe how burst activity in hippocampal place cells during navigation encodes spatial information. The burst frequency of place cells when animal is in field is around 8 Hz, suggesting that asynchronous release probably is important at neural circuit level.
 20. Stacey WC, Durand DM: **Stochastic resonance improves signal detection in hippocampal CA1 neurons.** *J Neurophysiol* 2000, **83**:1394-1402.
 21. Yu CR, Power J, Barnea G, O'Donnell S, Brown HE, Osborne J, Axel R, Gogos JA: **Spontaneous neural activity is required for the establishment and maintenance of the olfactory sensory map.** *Neuron* 2004, **42**:553-566.
 22. Harata N, Pyle JL, Aravanis AM, Mozhayeva M, Kavalali ET, Tsien RW: **Limited numbers of recycling vesicles in small CNS nerve terminals: implications for neural signaling and vesicular cycling.** *Trends Neurosci* 2001, **24**:637-643.
 23. Rizzoli SO, Betz WJ: **The structural organization of the readily releasable pool of synaptic vesicles.** *Science* 2004, **303**:2037-2039.
 24. Hubbard JI, Jones SF, Landau EM: **An examination of the effects of osmotic pressure changes upon transmitter release from mammalian motor nerve terminals.** *J Physiol* 1968, **197**:639-657.
 25. Rosenmund C, Stevens CF: **Definition of the readily releasable pool of vesicles at hippocampal synapses.** *Neuron* 1996, **16**:1197-1207.
 26. Xu-Friedman MA, Harris KM, Regehr WG: **Three-dimensional comparison of ultrastructural characteristics at depressing and facilitating synapses onto cerebellar Purkinje cells.** *J Neurosci* 2001, **21**:6666-6672.
 27. Gerber SH, Rah JC, Min SW, Liu X, de Wit H, Dulubova I, Meyer AC, Rizo J, Arancillo M, Hammer RE *et al.*: **Conformational switch of syntaxin-1 controls synaptic vesicle fusion.** *Science* 2008, **321**:1507-1510.
In this paper, mice in which syntaxin-1 is constitutively 'open' and thus activated are used to examine the relationship of SNARE-complex assembly to Ca²⁺-triggered vesicle exocytosis. The result identifies a function for the closed conformation of syntaxin-1 in controlling fusion of vesicles in the RRP via interaction with SM proteins, and demonstrates that vesicle docking is not equivalent to formation of the RRP.
 28. Voets T, Neher E, Moser T: **Mechanisms underlying phasic and sustained secretion in chromaffin cells from mouse adrenal slices.** *Neuron* 1999, **23**:607-615.
 29. Braun M, Ramracheya R, Johnson PR, Rorsman P: **Exocytotic properties of human pancreatic beta-cells.** *Ann N Y Acad Sci* 2009, **1152**:187-193.
 30. Lefkowitz JJ, Fogarty KE, Lifshitz LM, Bellve KD, Tuft RA, ZhuGe R, Walsh JV Jr, De Crescenzo V: **Suppression of Ca²⁺ syntillas increases spontaneous exocytosis in mouse adrenal chromaffin cells.** *J Gen Physiol* 2009, **134**:267-280.
 31. Li C, Ullrich B, Zhang ZZ, Anderson RGW, Brose N, Südhof TC: **Ca²⁺-dependent and Ca²⁺-independent activities of neural and nonneural synaptotagmins.** *Nature* 1995, **375**:594-599.

32. Sugita S, Shin O-H, Han W, Lao Y, Südhof TC: **Synaptotagmins form hierarchy of exocytotic Ca²⁺-sensors with distinct Ca²⁺-affinities.** *EMBO J* 2002, **21**:270-280.
33. Bhalla A, Tucker WC, Chapman ER: **Synaptotagmin isoforms couple distinct ranges of Ca²⁺, Ba²⁺, and Sr²⁺ concentration to SNARE-mediated membrane fusion.** *Mol Biol Cell* 2005, **16**:4755-4764.
34. Südhof TC, Rothman JE: **Membrane fusion: grappling with SNARE and SM proteins.** *Science* 2009, **323**:474-477.
35. Davletov BA, Südhof TC: **A single C₂-domain from synaptotagmin I is sufficient for high affinity Ca²⁺/phospholipid-binding.** *J Biol Chem* 1993, **268**:26386-26390.
36. Giraud CG, Garcia-Diaz A, Eng WS, Chen Y, Hendrickson WA, ●● Melia TJ, Rothman JE: **Alternative zippering as an on-off switch for SNARE-mediated fusion.** *Science* 2009, **323**:512-516.
This article reports a simple mechanistic model that accounts for the clamping function of complexin, a model whereby the accessory α -helix of complexin inserts into the assembling, partially zippered SNARE complex to prevent its full assembly, and is displaced from this position by Syt1.
37. Maximov A, Tang J, Yang X, Pang ZP, Südhof TC: **Complexin ●● controls the force transfer from SNARE complexes to membranes in fusion.** *Science* 2009, **323**:516-521.
Use genetics, RNAi knockdowns and electrophysiology, this study demonstrates that complexin functions in synaptic vesicle exocytosis by binding to assembling SNARE complexes and not individual SNARE proteins, and that this binding mediates both the clamping and activation functions of complex via distinct domains.
38. Tang J, Maximov A, Shin O-H, Dai H, Rizo J, Südhof TC: **A complexin/synaptotagmin-1 switch controls fast synaptic vesicle exocytosis.** *Cell* 2006, **126**:1175-1187.
39. Mackler JM, Drummond JA, Loewen CA, Robinson IM, Reist NE: **The C(2)B Ca(2+)-binding motif of synaptotagmin is required for synaptic transmission in vivo.** *Nature* 2002, **418**:340-344.
40. Shin OH, Xu J, Rizo J, Südhof TC: **Differential but convergent ● functions of Ca²⁺ binding to synaptotagmin-1 C2 domains mediate neurotransmitter release.** *Proc Natl Acad Sci USA* 2009, **106**:16469-16474.
This study demonstrates that blocking Ca²⁺-binding to the C2A-domain of Syt1 changes the apparent Ca²⁺-cooperativity of exocytosis, thereby providing formal proof that Ca²⁺-binding to both C2-domains of Syt1 contributes to triggering exocytosis.
41. Fernandez-Chacon R, Konigstorfer A, Gerber SH, Garcia J, Matos MF, Stevens CF, Brose N, Rizo J, Rosenmund C, Südhof TC: **Synaptotagmin I functions as a calcium regulator of release probability.** *Nature* 2001, **410**:41-49.
42. Pang ZP, Shin O-H, Meyer AC, Rosenmund C, Südhof TC: **A gain-of-function mutation in synaptotagmin-1 reveals a critical role of Ca²⁺-dependent SNARE-complex binding in synaptic exocytosis.** *J Neurosci* 2006, **26**:12556-12565.
43. Xu J, Mashimo T, Südhof TC: **Synaptotagmin-1, -2, and -9: ● Ca(2+) sensors for fast release that specify distinct presynaptic properties in subsets of neurons.** *Neuron* 2007, **54**:567-581.
This paper demonstrated that among all synaptotagmins, only Syt1, Syt2, and Syt9 can function as Ca²⁺-sensor for synaptic exocytosis in Syt1 knockout neurons, and additionally demonstrates that the three different synaptotagmin isoforms confer distinct properties onto Ca²⁺-triggered exocytosis.
44. Sun J, Pang ZP, Qin D, Fahim AT, Adachi R, Südhof TC: **A dual-●● Ca²⁺-sensor model for neurotransmitter release in a central synapse.** *Nature* 2007, **450**:676-682.
This paper describes the biophysical properties of asynchronous exocytosis using photolysis of caged Ca²⁺ and Ca²⁺-imaging in the calyx of Held synapse, and establishes the most precise model of Ca²⁺-triggered exocytosis up-to-date.
45. Voets T, Moser T, Lund PE, Chow RH, Geppert M, Südhof TC, Neher E: **Intracellular calcium dependence of large dense-core vesicle exocytosis in the absence of synaptotagmin I.** *Proc Natl Acad Sci USA* 2001, **98**:11680-11685.
46. Sorensen JB, Fernandez-Chacon R, Südhof TC, Neher E: **Examining synaptotagmin 1 function in dense core vesicle exocytosis under direct control of Ca²⁺.** *J Gen Physiol* 2003, **122**:265-276.
47. Li Y, Wang P, Xu J, Gorelick F, Yamazaki H, Andrews N, Desir GV: **Regulation of insulin secretion and GLUT4 trafficking by the calcium sensor synaptotagmin VII.** *Biochem Biophys Res Commun* 2007, **362**:658-664.
48. Schonn JS, Maximov A, Lao Y, Südhof TC, Sorensen JB: ● **Synaptotagmin-1 and -7 are functionally overlapping Ca²⁺ sensors for exocytosis in adrenal chromaffin cells.** *Proc Natl Acad Sci USA* 2008, **105**:3998-4003.
The data presented in this paper indicate that Syt1 and Syt7 together mediate nearly all Ca²⁺-triggered exocytosis in chromaffin cells, but execute different phases of vesicle exocytosis.
49. Gustavsson N, Lao Y, Maximov A, Chuang JC, Kostromina E, ● Repa JJ, Li C, Radda GK, Südhof TC, Han W: **Impaired insulin secretion and glucose intolerance in synaptotagmin-7 null mutant mice.** *Proc Natl Acad Sci USA* 2008, **105**:3992-3997.
Using mouse genetics, the authors not only show that Syt7 functions as a Ca²⁺-sensor in insulin secreting β -cells, but also demonstrate that another Ca²⁺-sensor must exist to account for all of the Ca²⁺-triggered exocytosis observed.
50. Gustavsson N, Wei SH, Hoang DN, Lao Y, Zhang Q, Radda GK, ● Rorsman P, Südhof TC, Han W: **Synaptotagmin-7 is a principal Ca²⁺ sensor for Ca²⁺-induced glucagon exocytosis in pancreas.** *J Physiol* 2009, **587**:1169-1178.
This study demonstrates that Syt7 functions as the major Ca²⁺-sensor for glucagon secretion in pancreatic α -cells.
51. Nagy G, Kim JH, Pang ZP, Matti U, Rettig J, Südhof TC, Sorensen JB: **Different effects on fast exocytosis induced by synaptotagmin 1 and 2 isoforms and abundance but not by phosphorylation.** *J Neurosci* 2006, **26**:632-643.
52. Maximov A, Lao Y, Li H, Chen X, Rizo J, Sorensen JB, Südhof TC: **Genetic analysis of synaptotagmin-7 function in synaptic vesicle exocytosis.** *Proc Natl Acad Sci USA* 2008, **105**:3986-3991.
53. de Wit H, Walter AM, Milosevic I, Gulyas-Kovacs A, Riedel D, ● Sorensen JB, Verhage M: **Synaptotagmin-1 docks secretory vesicles to syntaxin-1/SNAP-25 acceptor complexes.** *Cell* 2009, **138**:935-946.
The authors suggest that different from synapses, adrenal chromaffin cells use a complex of Syt1 with SNARE proteins and Munc18-1 for vesicle docking.
54. Melicoff E, Sansores-Garcia L, Gomez A, Moreira DC, Datta P, ● Thakur P, Petrova Y, Siddiqi T, Murthy JN, Dickey BF *et al.*: **Synaptotagmin-2 controls regulated exocytosis but not other secretory responses of mast cells.** *J Biol Chem* 2009, **284**:19445-19451.
This paper provides the best evidence available for the function of synaptotagmins as Ca²⁺-sensors for exocytosis in non-neuronal and non-endocrine cells, by demonstrating that Syt2 mediates Ca²⁺-induced mast cell exocytosis.
55. Martinez I, Chakrabarti S, Hellevik T, Morehead J, Fowler K, Andrews NW: **Synaptotagmin VII regulates Ca(2+)-dependent exocytosis of lysosomes in fibroblasts.** *J Cell Biol* 2000, **148**:1141-1149.
56. Jaiswal JK, Chakrabarti S, Andrews NW, Simon SM: **Synaptotagmin VII restricts fusion pore expansion during lysosomal exocytosis.** *PLoS Biol* 2004, **2**:E233.
57. Bollmann JH, Sakmann B, Borst JG: **Calcium sensitivity of glutamate release in a calyx-type terminal.** *Science* 2000, **289**:953-957.
58. Schneggenburger R, Neher E: **Intracellular calcium dependence of transmitter release rates at a fast central synapse.** *Nature* 2000, **406**:889-893.
59. Meinrenken CJ, Borst JG, Sakmann B: **Local routes revisited: the space and time dependence of the Ca²⁺ signal for phasic transmitter release at the rat calyx of Held.** *J Physiol* 2003, **547**:665-689.
60. Franks KM, Stevens CF, Sejnowski TJ: **Independent sources of quantal variability at single glutamatergic synapses.** *J Neurosci* 2003, **23**:3186-3195.

61. Groffen AJ, Martens S, Arazola RD, Cornelisse LN, Lozovaya N, de Jong AP, Goriounova NA, Habets RL, Takai Y, Borst JG *et al.*: **Doc2b is a high-affinity Ca²⁺ sensor for spontaneous neurotransmitter release.** *Science* 2010, **327**:1614-1618.
- This paper reports that Doc2B, a cytosolic double C2 domain containing protein with higher Ca²⁺-affinity than Syt1, associates with phospholipids in the presence of Ca²⁺, binds to SNARE complexes, and promotes membrane fusion. *In vivo* data suggest that Doc2B partly contributes to spontaneous synaptic vesicle exocytosis.
62. Sara Y, Virmani T, Deak F, Liu X, Kavalali ET: **An isolated pool of vesicles recycles at rest and drives spontaneous neurotransmission.** *Neuron* 2005, **45**:563-573.
63. Fredj NB, Burrone J: **A resting pool of vesicles is responsible for spontaneous vesicle fusion at the synapse.** *Nat Neurosci* 2009, **12**:751-758.
- This study used a novel technique to demonstrate that synaptic vesicles responsible for spontaneous release are distinct from synaptic vesicles that are released in response to action potentials.
64. Pang ZP, Cao P, Xu W, Südhof TC: **Calmodulin controls synaptic strength via presynaptic activation of CaM kinase II.** *J Neurosci* 2010, **30**:4132-4442.
65. Washington NL, Ward S: **FER-1 regulates Ca²⁺-mediated membrane fusion during *C. elegans* spermatogenesis.** *J Cell Sci* 2006, **119**:2552-2562.
66. Bansal D, Miyake K, Vogel SS, Groh S, Chen CC, Williamson R, McNeil PL, Campbell KP: **Defective membrane repair in dysferlin-deficient muscular dystrophy.** *Nature* 2003, **423**:168-172.
67. Roux I, Safieddine S, Nouvian R, Grati M, Simmler MC, Bahloul A, Perfettini I, Le Gall M, Rostaing P, Hamard G *et al.*: **Otoferlin, defective in a human deafness form, is essential for exocytosis at the auditory ribbon synapse.** *Cell* 2006, **127**:277-289.
68. Shin OH, Han W, Wang Y, Südhof TC: **Evolutionarily conserved multiple C2 domain proteins with two transmembrane regions (MCTPs) and unusual Ca²⁺ binding properties.** *J Biol Chem* 2005, **280**:1641-1651.
69. Min SW, Chang WP, Südhof TC: **E-Syts, a family of membranous Ca²⁺-sensor proteins with multiple C2 domains.** *Proc Natl Acad Sci USA* 2007, **104**:3823-3828.
70. Deak F, Shin OH, Tang J, Hanson P, Ubach J, Jahn R, Rizo J, Kavalali ET, Südhof TC: **Rabphilin regulates SNARE-dependent re-priming of synaptic vesicles for fusion.** *EMBO J* 2006, **25**:2856-2866.
71. Friedrich R, Groffen AJ, Connell E, van Weering JR, Gutman O, Henis YI, Davletov B, Ashery U: **DOC2B acts as a calcium switch and enhances vesicle fusion.** *J Neurosci* 2008, **28**:6794-6806.
- This paper utilizes chromaffin cells as a model system to study the function of Doc2B, showing that Doc2B functions as a priming factor and controls the fusogenicity of LDCVs.
72. Tomsig JL, Creutz CE: **Copines: a ubiquitous family of Ca(2+)-dependent phospholipid-binding proteins.** *Cell Mol Life Sci* 2002, **59**:1467-1477.
73. Yoshihara M, Adolfsen B, Galle KT, Littleton JT: **Retrograde signaling by Syt 4 induces presynaptic release and synapse-specific growth.** *Science* 2005, **310**:858-863.
74. Südhof TC: **Synaptotagmins: why so many?** *J Biol Chem* 2002, **277**:7629-7632.
75. Maximov A, Shin OH, Liu X, Südhof TC: **Synaptotagmin-12, a synaptic vesicle phosphoprotein that modulates spontaneous neurotransmitter release.** *J Cell Biol* 2007, **176**:113-124.
76. Gustavsson N, Han W: **Calcium-sensing beyond neurotransmitters: functions of synaptotagmins in neuroendocrine and endocrine secretion.** *Biosci Rep* 2009, **29**:245-259.
77. Choi UB, Strop P, Vrljic M, Chu S, Brunger AT, Weninger KR: **Single-molecule FRET-derived model of the synaptotagmin 1-SNARE fusion complex.** *Nat Struct Mol Biol* 2010, **17**:318-324.
78. Vrljic M, Strop P, Ernst JA, Sutton RB, Chu S, Brunger AT: **Molecular mechanism of the synaptotagmin-SNARE interaction in Ca²⁺-triggered vesicle fusion.** *Nat Struct Mol Biol* 2010, **17**:325-331.
79. Lee HK, Yang Y, Su Z, Hyeon C, Lee TS, Lee HW, Kweon DH, Shin YK, Yoon TY: **Dynamic Ca²⁺-dependent stimulation of vesicle fusion by membrane-anchored synaptotagmin 1.** *Science* 2010, **328**:760-763.